

Review

Nitroxidative Signaling Mechanisms in Pathological Pain

Peter M. Grace,^{1,4,*} Andrew D. Gaudet,¹ Vasiliki Staikopoulos,² Steven F. Maier,¹ Mark R. Hutchinson,² Daniela Salvemini,³ and Linda R. Watkins¹

Tissue injury can initiate bidirectional signaling between neurons, glia, and immune cells that creates and amplifies pain. While the ability for neurotransmitters, neuropeptides, and cytokines to initiate and maintain pain has been extensively studied, recent work has identified a key role for reactive oxygen and nitrogen species (ROS/RNS; nitroxidative species), including superoxide, peroxynitrite, and hydrogen peroxide. In this review we describe how nitroxidative species are generated after tissue injury and the mechanisms by which they enhance neuroexcitability in pain pathways. Finally, we discuss potential therapeutic strategies for normalizing nitroxidative signaling, which may also enhance opioid analgesia, to help to alleviate the enormous burden of pathological pain.

The Link Between Nitroxidative Signaling and Pain

Investigation of oxidative processes, such as rusting, began with the ‘phlogiston theory’, developed by Georg Ernest Stahl during the scientific revolution, which postulated that a fire-like element (phlogiston) is released during combustion. Oxidation was formally linked to biology during the early 20th Century, when it was found to underpin cellular metabolism [1–3]. The connection between ROS and altered sensory processing was empirically identified around the same time [4]. Since then, research has shown that prolonged, unchecked increases in ROS/RNS (nitroxidative species) after infection or tissue damage can promote cytotoxicity and inflammation. These processes can cause peripheral and central sensitization which underlie **pathological pain** (see *Glossary*) [5,6]. Thus, restoring nitroxidative balance in peripheral and central nervous systems (PNS, CNS) is a possible therapeutic approach for ameliorating neuropathology [6–10].

In this review, we summarize recent research on how nitroxidative species participate in **neuroimmune signaling** throughout the neuraxis to drive pathological pain. We additionally discuss potential therapeutic strategies for normalizing nitroxidative signaling by activating endogenous antioxidant systems, which may also enhance opioid analgesia. Because pathological pain is often intractable to current therapies, new strategies to normalize nitroxidative signaling may help to alleviate the enormous burden of pain [11].

Production of Nitroxidative Species by Neurons, Glia, and Immune Cells

The role of nitroxidative signaling in pain has been studied using rodent experimental models of inflammatory pain [e.g., intraplantar complete Freund's adjuvant (CFA), formalin] and neuropathic pain [e.g., peripheral nerve injury (PNI), chemotherapy-induced peripheral neuropathy

Trends

Nitroxidative species (RNS, ROS, and their products) contribute to peripheral and central sensitization after tissue injury, which leads to pathological pain.

There is a reciprocal relationship between nitroxidative and inflammatory signaling that drives peripheral and central sensitization.

New approaches to restoring nitroxidative balance may reveal effective strategies to treat pathological pain.

The development of new tools may enhance our understanding of the crucial relationships between real-time local creation of nitroxidative species, their concentration at the effect site, and the distribution of their direct effects.

¹Department of Psychology and Neuroscience, and the Center for Neuroscience, University of Colorado, Boulder, CO, USA

²Discipline of Physiology, School of Medicine, and the Australian Research Council (ARC) Centre for Nanoscale BioPhotonics (CNBP), University of Adelaide, Adelaide, SA, Australia

³Department of Pharmacology and Physiology, Saint Louis University School of Medicine, St. Louis, MO, USA

⁴Current address: Department of Critical Care Research, University of Texas MD Anderson Cancer Center, Houston, TX, USA

*Correspondence: pgrace@mdanderson.org (P.M. Grace).




Figure 1. Induction of Nitroxidative Species after Tissue Injury. Nitroxidative species can induce post-translational modifications of proteins and lipids, which subsequently drive pathological pain by modulating nociceptive neurotransmission, activating TRP channels, and inducing mitochondrial dysfunction, and thereby induce inflammatory signaling. In healthy cells, endogenous antioxidant systems prevent nitroxidative damage. Cell damage/pathology can perturb this balance, driving the accumulation of potentially damaging nitroxidative species. Abbreviations: CAT, catalase; GPx, glutathione; H₂O₂, hydrogen peroxide; HO, heme oxygenase; mETC, mitochondrial electron transport chain; NO, nitric oxide; NOS, nitric oxide synthase; NOX, NADPH oxidase; O₂^{•-}, superoxide; •OH, hydroxyl radical; ONOO⁻, peroxynitrite; SOD, superoxide dismutase.

Glossary

Neuroimmune signaling: bidirectional communication between leukocytes, glia, and neurons.

Pathological pain: maladaptive pain that serves no useful purpose.

(CIPN), diabetic neuropathy (DN), spinal cord injury (SCI), experimental autoimmune encephalomyelitis (EAE)], which have recently been reviewed elsewhere [12]. There are numerous endogenous sources of ROS and nitric oxide (NO) that are engaged during pain processing [13]. NADPH oxidases, NO synthases, and mitochondrial respiration are among the best characterized ROS/NO producers, and will be discussed here (Figures 1 and 2).

NADPH Oxidases

NADPH oxidases (NOX) are membrane-bound enzyme complexes. They transport electrons donated from cytosolic NADPH to generate extracellular or luminal superoxide anions or hydrogen peroxide that can be transported into the cytosol via aquaporin channels [13,14]. In contrast to other sources of ROS that are generated as a byproduct of catabolism, ROS generation is the primary function of NOX. There are seven members in the NOX family; NOX1, 2, and 4 have been implicated in pathological inflammatory and neuropathic pain models [13,15,16]. NOX1 and 2 are expressed at the cellular membrane, and produce superoxide anions following phosphorylation of cytosolic subunits [17]. NOX4 is expressed on organelles, such as the endoplasmic reticulum, and constitutively produces hydrogen peroxide [17].

NOX1 is inducibly expressed by microglia, neurons, astrocytes, and macrophages in the dorsal root ganglion (DRG) and CNS [17–19]. Nociceptive hypersensitivity induced by the inflammatory stimuli formalin and carrageenan is attenuated in *Nox1*-deficient mice [18]. NOX1-derived ROS induce translocation of PKC ϵ to the membrane to enhance transient receptor potential (TRP) V1 activity in DRG neurons [18], a change consistent with pain amplification (Figure 2). By contrast, another study showed that NOX1 mRNA failed to upregulate in the DRG following peripheral nerve injury (PNI) [20]. These results indicate that DRG NOX1 may have a preferential role in inflammatory versus neuropathic pain.

NOX2 is predominantly expressed by phagocytic cells – peripheral macrophages and CNS microglia [13]. PNI induces a rapid upregulation of NOX2 mRNA by DRG macrophages and spinal microglia, which is correlated with increased intracellular superoxide [20,21]. PNI-induced nociceptive hypersensitivity was attenuated in *Nox2*-deficient mice [20,21]. *Nox2* deficiency attenuated TNF, but not IL-1 β , mRNA expression, as well as expression of the neuronal injury marker ATF3 in DRG (Figure 2) [20]. However, *Nox2* deficiency did not influence macrophage recruitment to the injured DRG, suggesting a role for NOX2 in macrophage function rather than in




Figure 2. Sources of Nitroxidative Species after Tissue Injury. Principal sources of nitroxidative species include NADPH oxidase (NOX), nitric oxide synthase (NOS), and electron leakage from the mitochondrial electron transport chain (mETC). The NOX1, 2, and 4 isoforms are differentially expressed across cell types and tissues after injury. NOX1-derived reactive oxygen species (ROS) induce enhanced transient receptor potential (TRP) V1 activity in dorsal root ganglia (DRG) neurons. NOX2 activity in macrophages and microglia drives proinflammatory cytokine (PIC) gene expression in DRG of the spinal dorsal horn. NOX4 expression at the site of peripheral nerve injury decreases the expression of myelin proteins (MP). The three NOS isoforms – NOS1 (neuronal), 2 (inducible), and 3 (endothelial) – are also differentially expressed by cell type. In abnormal pain states, *N*-methyl-D-aspartate receptors (NMDARs) are activated, resulting in calcium influx and activation of NOS1. Transcription of NOS2 is initiated by Toll-like receptors (TLRs). These enzymes and processes have a well-established role in pathological pain.

chemotaxis [20]. *Nox2* deficiency attenuated PNI-induced *Iba1* expression and the attendant expression of proinflammatory cytokines TNF and IL-1 β in the spinal dorsal horn [21]. Because these studies were performed in global knockouts, it is still unclear whether alterations in the DRG and dorsal horn are subject to NOX-dependent changes in macrophage function at the injury site. In contrast to NOX1, NOX2 activity in monocytes appears to play no role in inflammatory pain [22].

NOX4 is expressed by DRG neurons – both myelinated (A-fibers) and unmyelinated (C-fibers) DRG neurons – as well as by microglia, astrocytes, and macrophages [13,23,24]. Nociceptive hypersensitivity following PNI is attenuated in *Nox4*-deficient mice, with attenuation of hydrogen peroxide at the sciatic nerve injury site [23]. These results are supported by the absence of NOX4 upregulation in the DRG after PNI [20]. The myelin proteins MPZ and PMP22 are decreased at

the sciatic nerve injury site over time in a NOX4-dependent fashion, suggesting that myelin degeneration by hydrogen peroxide may maintain neuropathic pain (Figure 2). However, attenuated damage at the injury site did not alter expression of the nitroxidative stress and neuroinflammation indices at the spinal dorsal horn or DRG (microglia proliferation, hydrogen peroxide levels) [23]. This contrasts with other studies showing that such processes are dependent on manipulations at the sciatic nerve [25–27]. Finally, a role for NOX4 may be limited to neuropathic rather than inflammatory pain [23].

Together, these data suggest that NOX1, 2, and 4 isoforms contribute to pathological pain. Future studies could expand the role of various NOX isoforms to other sites in the neuraxis and well as identifying a role for other NOX isoforms in pain. These data may help to guide the development of therapeutics that target the activity of specific NOX isoforms to reduce nitroxidative stress and pain.

Nitric Oxide Synthases

NO is a diffusible gas mediator that is synthesized from L-arginine by one of three nitric oxide synthase (NOS) isoforms: NOS1 (neuronal), 2 (inducible), and 3 (endothelial). NO and all three NOS isoforms have a well-established role in nociception (Figure 2) [28]. It easily passes through membranes to directly impact on nearby cells.

NOS1 is constitutively expressed in the cytosolic compartment of postsynaptic terminals of neurons, and of stressed Schwann cells, and requires calcium for its activation [29–31]. In abnormal pain states, *N*-methyl-D-aspartate (NMDA) receptors are activated, resulting in calcium influx and activation of NOS1 [28]. Nociceptive hypersensitivity induced by PNI and CIPN is attenuated by genetic ablation and pharmacological inhibition of NOS1 [32–35].

NOS2 is a cytosolic isoform that is widely expressed in many immune cells and in glia. Transcription of NOS2 is initiated by Toll-like receptors (TLRs) and, once translated, is constitutively active – that is, unlike NOS1 and 3, its activity is independent of calcium [28]. NOS2 inhibition attenuates nociceptive hypersensitivity associated with inflammatory and neuropathic pain models [15,36,37].

NOS3 is best known for its expression in the cardiovascular system as a regulator of vascular tone. NOS3 is a membrane-bound enzyme that is constitutively expressed; however, it requires the interaction of calcium and calmodulin for its activation [28]. NOS3 expression is increased in the DRG after subcutaneous administration of CFA, and is correlated with allodynia, suggestive of increased NOS3 activity [38]. CFA-induced inflammatory pain is attenuated by NOS3 inhibition [38].

Cellular Respiration

One crucial function of mitochondria is in energy metabolism. The mitochondrial electron transport chain (mETC) is a series of five molecular complexes through which electrons are transported to synthesize ATP from ADP. Premature electron leakage can occur during cellular respiration, particularly at complexes I and III, resulting in superoxide production (Figure 2) [39].

Mitochondrial ROS levels are elevated in spinal neurons, microglia, and astrocytes in neuropathic pain models [21,40,41]. Furthermore, blocking the mETC attenuates hyperalgesia associated with a range of inflammatory and neuropathic pain models [42–45]. However, a direct link between mETC-dependent pain and mitochondrial ROS has yet to be shown. These results suggest that cellular respiration is increased, but is inefficient owing to enhanced ROS-generating electron leakage from the mETC because ATP production by sciatic nerves is impaired during CIPN [46].

Mechanisms of Nitroxidative Signaling in Neuronal Hyperexcitability

Injury or disease can provoke intense, repeated, and sustained activity of primary afferent (sensory) neurons. This activity, together with the release of mediators from reactive glia and immune cells, elicits well-characterized changes in neuronal and biochemical processing at peripheral terminals and central synapses [5,47–50]. This is termed ‘sensitization’, and results in nociceptive hypersensitivity. We discuss here how nitroxidative signaling engages neurons in pain pathways, leading to peripheral and central sensitization (Figures 1 and 3).

Nitroxidative Species as Neuromodulators in Pain Pathways


Nitroxidative species can directly increase the excitability of nociceptive neurons. Intraplantar administration of superoxide or peroxynitrite, or intrathecal delivery of the ROS donor *tert*-butyl hydroperoxide (tBOOH), is sufficient to induce nociceptive hypersensitivity in naïve rats [51–54]. These studies demonstrated that ROS activates calcium/calmodulin-dependent protein kinase II (CaMKII) in glutamatergic spinal neurons, and induced presynaptic inhibition of GABAergic interneurons (disinhibition). Furthermore, hydrogen peroxide enhanced the frequency and amplitude of action potentials of DRG neurons from neuropathic rats (Figure 3) [55].

In neuropathic pain models, administration of the non-selective ROS scavenger phenyl-*N*-*tert*-butylnitronate (PBN), selective small-molecule superoxide and peroxynitrite decomposition catalysts such as M40403, FeTMAPyP⁵⁺, and MnTE-2-PyP⁵⁺, or selective peroxynitrite decomposition catalysts such as SRI6 and SRI110, attenuated nociceptive hypersensitivity [15,51,53,54,56–58]. Accordingly, PBN attenuated injury-induced hyperexcitability of spinal dorsal (sensory) horn ‘pain’-responsive neurons and phosphorylation of CaMKII [51,57], an effect consistent with pain normalization. Several mechanisms of enhanced excitatory signaling have been identified. Hydrogen peroxide can activate cGK α , resulting in increased neurotransmitter release from the terminals of primary afferent neurons in the dorsal horn [59,60]. Peroxynitrite and ROS disrupt glutamate homeostasis, leading to potentiation of synaptic currents and calcium influx, and ultimately excitotoxicity [56,61]. Mechanisms include nitration and phosphorylation of several NMDA receptor subunits, as well as inhibition of glutamine synthetase and the glutamate transporter GLT-1 that limit the synaptic half-life of glutamate [15,56,62,63]. Nitroxidative products also induce disinhibition after PNI because PBN normalized the decrease in GAD-67⁺ GABAergic dorsal horn neurons, and increased GABA release (Figure 3) [53,64]. Together, these data suggest that nitroxidative species directly enhance neuroexcitability in pain pathways.

Nitroxidative Species Activate TRP Channels

The TRP family of nonselective cation channels play a vital role in the molecular integration of multiple endogenous and exogenous sensory stimuli [65]. Several of these channels, expressed at the peripheral and central terminals and cell bodies of primary afferent neurons, are activated by nitroxidative species and products. TRP channel activation by nitroxidative species can also initiate neurogenic inflammation – recruitment and activation of immune cells following release of neuropeptides by neurons – which is a key process underlying pathological pain (Figure 3) [5,66]. We focus here on the known roles of TRPA1, TRPM2, and TRPV1.

TRPA1 is a chemoreceptor expressed exclusively by peptidergic C-fibers [65]. Nitroxidative species induce protein carbonylation, and membrane phospholipid peroxidation and nitration, as well as subsequent production of reactive aldehydes such as acrolein (Figure 1). These products all share the ability to induce nociceptive hypersensitivity by directly activating TRPA1 [67–72]. Acrolein is elevated in the DRG and spinal cord after SCI, and blockade with hydralazine or phenelzine partially attenuated allodynia [73,74]. Moreover, nociceptive hypersensitivity induced by CIPN was abolished in *Trpa1*-deficient mice or by a TRPA1 antagonist [75]. In this



Trends in Neurosciences

Figure 3. Nitroxidative Mechanisms of Neuroexcitability after Tissue Injury. Reactive nitroxidative species, such as hydrogen peroxide and peroxynitrite, and modified proteins and lipids, including carbonylated proteins, peroxidized and nitrated lipids, and reactive aldehydes, all contribute to peripheral and central sensitization after tissue injury. These processes drive pathological pain. Several of the transient receptor potential (TRP) family of nonselective cation channels are activated by nitroxidative species and by modified proteins and lipids (see Nitroxidative Species Activate TRP Channels). TRPA1 is expressed by peptidergic C-fibers, and is activated by modified proteins and lipids. TRPM2, which is expressed by neurons, monocytes/macrophages, microglia, and T cells, is directly activated by nitroxidative species. TRPM2 also activates intracellular signaling pathways, including mitogen-activated protein kinase (MAPK) pathways and nuclear translocation of nuclear factor κ light-chain enhancer of activated B cells (NF- κ B). TRPV1 is found on C-fibers and is directly activated by some modified proteins and lipids, as well as being a target of oxidation and nitration events by nitroxidative species that increase the responsiveness of the channel. Reactive nitroxidative species can directly modulate neuroexcitability in central synapses by promoting glutamate release from primary afferent terminals, by activating calcium/calmodulin-dependent protein kinase II (CaMKII) in glutamatergic spinal neurons, and by inhibiting GABAergic interneurons (see Nitroxidative Species as Neuromodulators in Pain Pathways). Nitroxidative species also disrupt glutamate homeostasis by nitration and phosphorylation of NMDA receptor (NMDAR) subunits, as well as by inhibiting glutamine synthetase (GS) and the glutamate transporter GLT-1. Mitochondrial DNA is a target for oxidation and nitration, while some nitroxidative species can form adducts with many mitochondrial proteins, which together impair the structural integrity and function of mitochondria (see Nitroxidative Species Induce Mitochondrial Dysfunction). Nitroxidative species can also trigger release of proapoptotic factors from mitochondria by disrupting organelle dynamics. Nitroxidative species induce the production of proinflammatory mediators, and can activate NF- κ B and MAPK intracellular signaling pathways (see Nitroxidative Species Induce Neuroinflammatory Signaling). Toll-like receptors (TLRs) bind to a variety of endogenous danger signals, including those released from nitroxidative-damaged mitochondria, to activate NF- κ B and MAPKs. NOX-derived ROS are second

(Figure legend continued on the bottom of the next page.)

model, the chemotherapeutic bortezomib did not directly activate TRPA1, suggesting that ROS may act as an intermediate [75].

TRPM2 is expressed by neurons, and abundantly by immune cells including monocytes/macrophages, neutrophils, T cells, and microglia. This channel is directly activated by hydrogen peroxide and by cytosolic ADP-ribose that is generated after nitroxidative damage to mitochondria [76–81]. Furthermore, TRPM2 activation is crucial for activation of spinal microglia and for macrophage infiltration into the spinal cord after PNI [82]. TRPM2 also activates the ERK/MAPK pathway and induces nuclear translocation of NF- κ B, resulting in the production of proinflammatory cytokines and chemokines [76,77,81,83,84]. Consequently, pharmacological and genetic studies have demonstrated that TRPM2 contributes to inflammatory and neuropathic nociceptive hypersensitivity [77–79,82,85].

TRPV1 is found on unmyelinated, slowly-conducting neuronal C-fibers, and is an essential component underlying injury-elicited thermal hyperalgesia and nociceptive hypersensitivity [65]. TRPV1 expression is upregulated by an exogenous ROS donor (tBOOH), and is a target of oxidation and nitration events that increase the responsiveness of the channel [18,86–88]. Moreover, linoleic acid metabolites, created during the production of eicosanoids, are endogenous TRPV1 agonists when oxidized, and contribute to nociceptive signaling [89,90].

Nitroxidative Species Induce Mitochondrial Dysfunction

Mitochondria have pivotal roles in a variety of cellular functions, including energy metabolism, calcium homeostasis, lipid synthesis, and apoptosis. As noted above, cellular respiration can be elevated under neuropathic pain conditions, with an attendant elevation of ROS derived from neuronal and microglial mitochondria [21,40,41]. Together with nitroxidative species derived from NOX and NOS enzymes, these species disrupt mitochondrial homeostasis via several mechanisms, leading to bioenergetic crisis (due to impaired mETC efficiency) and degeneration of primary afferents (Figure 3) [91].

Mitochondrial DNA is a target of oxidation and nitration, while peroxidated lipid end-products, such as reactive aldehydes, can form covalent modifications (adducts) with an array of mitochondrial proteins, including antioxidants [92,93]. Together, these changes impair the structural integrity and function of mitochondria. Nitroxidative species can also trigger release of proapoptotic factors from mitochondria. For example, NO can disrupt mitochondrial dynamics (fission and fusion; responsible for maintaining metabolic homeostasis), resulting in translocation of Bcl-2-associated X protein from the cytosol to the organelle membrane, where it activates apoptosis pathways [94–96]. Activation of apoptosis pathways contributes to neuropathic pain because inhibition of several caspase enzymes attenuates vincristine- and dideoxycytidine-induced nociceptive hypersensitivity [97]. Neuropathic pain is associated with impaired mitochondrial function, and nociceptive hypersensitivity is accordingly attenuated by pharmacologically normalizing mitochondrial dynamics or preventing mitotoxicity [46,98–100].

Nitroxidative Species Induce Neuroinflammatory Signaling

Proinflammatory mediators released by glial and immune cells increase neuroexcitability in pain pathways after injury (e.g., TNF, IL-1 β , BDNF) [5,50,101–103]. Several mechanisms include enhanced glutamate release, increased AMPA receptor expression, phosphorylated NMDA receptor subunits, and downregulated astrocyte glutamate transporters [5]. These

messengers for NF- κ B- and p38 MAPK-dependent TLR signaling, and TLR expression. The TLR2–NOX1 interaction also upregulates adhesion molecules via CCL3, which facilitates transendothelial cell migration into the CNS. Mitochondria-derived ROS also activate NLRP3 inflammasomes, which are protein complexes responsible for the proteolytic activation of IL-1 β .

proinflammatory mediators can also induce disinhibition of neuronal excitability by attenuating GABA and glycine release from interneurons and inhibitory descending projections, and by downregulating KCC2 on postsynaptic terminals [5].

Nitroxidative species regulate the production of proinflammatory mediators during pathological pain. For example, NF- κ B and p38 MAPK are responsible for the production of a wide array of proinflammatory mediators in immune cells. Nitroxidative products degrade/inhibit I κ B and MAPK phosphatases, resulting in activation of NF- κ B and p38 that both mediate inflammatory and neuropathic pain [52,104–107]. Furthermore, nitroxidative species may promote the release of neuron-to-glia signals such as matrix metalloproteases (MMPs) (Figure 3) [108].

Nitroxidative species also elicit proinflammatory responses via TLR signaling. TLRs bind a variety of endogenous ligands (danger-associated molecular patterns, DAMPs), including DNA and N-formyl peptides from nitroxidatively damaged mitochondria, to trigger innate immune responses that contribute to pathological pain [5,109]. ROS serve a vital role as second messengers for TLR signaling. A rapid (minutes) respiratory burst occurs upon activation of TLR2 and 4, which is mediated by a direct interaction with the intracellular domains of NOX1, 2, and 4 enzymes. This NOX activity is essential for downstream NF- κ B- and p38 MAPK-dependent cytokine production [110–114]. Furthermore, activation of NOX enzymes by TLR signaling induces transcription of TLRs, and promotes membrane expression in lipid rafts, which is necessary for efficient signaling [111,115,116]. In concert with disruption of blood–brain barrier tight junctions by nitroxidative species, the TLR2–NOX1 interaction also upregulates adhesion molecules via CCL3 to facilitate transendothelial cell migration, which contributes to nociceptive hypersensitivity after PNI (Figure 3) [102,110,117].

ROS have been implicated in the activation of NLRP3 inflammasomes [118] – protein complexes responsible for the proteolytic activation of IL-1 β , a proinflammatory cytokine with a well-established role in pathological pain [5,101,119]. Among the various sensor molecules that trigger the formation of inflammasomes, NLRP3 has been most widely investigated and has a recently described role in neuropathic pain [120]. The relative contributions of ROS to the activation versus priming of NLRP3 inflammasomes remains to be elucidated [119]. Mitochondria are key participants in the activation of NLRP3 inflammasomes: they are a source of ROS that can directly activate NLRP3, as well as of oxidized mitochondrial DNA that can also activate NLRP3 (Figure 3) [118,121–123]. Furthermore, TRPM2 activation by nitroxidative species induces a calcium flux that activates the NLRP3 inflammasome [124].

Finally, there is a reciprocal relationship between nitroxidative species and inflammatory signaling. For example, the transcription of NOX and NOS enzymes is upregulated by TLR4 and 9 signaling, as well as by NF- κ B and p38 activation [19,125–129]. The purinergic receptor P2X7, which has a documented role in pathological pain, also induces ROS production [5,120,130]. ATP signaling through P2X7R activates NOX2 in a calcium and p38-dependent fashion [131–133].

Endogenous Regulators of Nitroxidative Signaling

Under healthy conditions, nitroxidative species and antioxidants exist in a balanced state because nitroxidative products play a vital physiological role in cellular processes (e.g., signal transduction, pathogen defense [134–136]). In response to increased production of nitroxidative species during injury or infection, antioxidant and regulatory systems are activated in an attempt to recover homeostasis (Figure 1) [14].

Antioxidant Defense

Transcription of antioxidant genes is a crucial step in controlling nitroxidative signaling. One key transcription factor is nuclear factor E2-related factor 2 (Nrf2). Nrf2 is expressed in CNS and PNS

neurons, macrophages, Schwann cells, astrocytes, and microglia [137–139]. Under homeostatic conditions, cytosolic Nrf2 is sequestered by the protein Keap1 and ubiquitinated for degradation. However, in the presence of oxidants and electrophiles, Nrf2 is released from Keap1 and translocates to the nucleus [140]. Nrf2 binds to the antioxidant response element (ARE) promoter region to elicit expression of >200 antioxidant genes, including those encoding superoxide dismutases (SOD1, cytosolic; SOD2, mitochondrial), catalase, glutathione, and heme-oxygenases [140]. Another transcription factor, forkhead box class O (FoxO), is also responsible for the production of SOD2 and catalase [141]. Many of these antioxidants are ubiquitously expressed, and their catabolic function is summarized in Figure 1 [142].

These endogenous antioxidant systems collaborate to detoxify reactive nitroxidative species (Figure 1). The evidence is mixed whether neuroinflammatory or traumatic events increase nervous system antioxidant levels [143–152]. This likely reflects a temporally- and injury-specific antioxidant response, and the fact that injury-induced nitroxidative species can negatively regulate antioxidant production [15,76]. Antioxidant system activation can limit pathological pain: deletion of SOD1 exacerbates neuropathic pain, while exogenous antioxidants attenuate nociceptive hypersensitivity in a range of inflammatory and neuropathic pain models [37,108,153–155]. Similarly, heme-oxygenases, which elicit the expression of various antioxidants, protect cells and could improve inflammation and neuropathic pain [21]. Therefore, therapies that increase antioxidant systems could resolve neuroinflammation and pain symptoms.

Anti-Inflammatory Cytokine and Adenosine Signaling

Cytokines such as IL-10 and TGF- β counter-regulate proinflammatory signaling and contribute to the resolution of neuropathic pain hypersensitivity [5,156,157]. One mechanism of action is via the regulation of nitroxidative signaling. For example, IL-10 and TGF- β inhibit NOX2 activity and promote antioxidant production [158–160]. This is a reciprocal relationship because antioxidants can also drive the production of anti-inflammatory cytokines [161,162]. Adenosine signaling is also anti-nociceptive in pathological pain models [163–165]. Signaling through A_{2A} and A₃ receptors inhibits NOX activity, and drives the production of anti-inflammatory cytokines and antioxidants [163,166,167].

Opposition of Opioid Analgesia by Nitroxidative Species

Opioid analgesics remain the cornerstone of management of moderate-to-severe pain. However, the clinical utility of opioids is limited by tolerance, which is characterized by dose escalation as a result of reduced sensitivity to an opioid agonist, as well as by hyperalgesia, a paradoxical increase in pain sensitivity induced by opioid exposure [168,169]. Recent evidence has identified a role for nitroxidative signaling in these phenomena [6,170].

NOX activity is elevated by morphine, and genetic or pharmacological disruption of these enzymes attenuates tolerance and hyperalgesia [171–173]. Superoxide and peroxynitrite have been implicated as downstream mediators because decomposition catalysts also attenuate tolerance and hyperalgesia [174–176]. It remains unclear how morphine engages these enzymes, but it may be mediated by classical μ -opioid receptors and/or TLR4 [168]. The pronociceptive mechanisms of nitroxidative species, described above, may act as an opposing process of neuronally-mediated opioid analgesia to create tolerance, or may overshadow analgesia to induce hyperalgesia. Therefore, correcting nitroxidative imbalance may improve the clinical profile of opioids [170].

Nitroxidative signaling also disrupts endogenous opioid analgesia in supraspinal sites that are engaged to inhibit spinal nociception via descending projections. For example, upregulation of peroxynitrite during inflammatory pain results in nitration of met-enkephalin in the rostral

ventromedial medulla (RVM), and this reduces opioid receptor binding affinity [177]. This may be normalized by intra-RVM microinjections of FeTMPyP⁵⁺, which was antinociceptive in inflammatory and neuropathic pain models [177].

Nitroxidative Signaling as a Therapeutic Target for Pathological Pain

Under pathological conditions, endogenous antioxidant responses can be insufficient, leading to an accumulation of toxic nitroxidative species. As mentioned above, unchecked increases in nitroxidative species can promote cytotoxicity and inflammation via cascading pronociceptive signaling. Therefore, discovering therapeutic treatments that enhance cellular antioxidant capacity could help to achieve nitroxidative balance to recover homeostasis.

Initial efforts to combat increases in nitroxidative species in a wide range of neurological disorders used direct antioxidant compounds (e.g., vitamins C and E, coenzyme Q). The consensus view is that the possible beneficial effects are outweighed by unfavorable pharmacokinetic and pharmacodynamic profiles [13,178,179]. A variety of redox-active therapeutics are being developed to overcome these issues, and are effective in treating cancer-induced bone pain as well as inflammatory and neuropathic pain, and can also potentiate opioid analgesia [9,10,180].

Newer approaches have instead aimed to inhibit sources of nitroxidative species, stimulate endogenous antioxidants, and prevent nitroxidative damage [13,178]. To this end, inhibitors of specific NOX and NOS isoforms, and ROS toxifiers such as MPO, are being developed and may prove effective for pain treatment [13,181]. As noted above, A_{2A} and A₃ adenosine receptor agonists attenuate spinal NOX activity and promote antioxidant production, with a concomitant decrease in neuropathic pain [163–165]. Another promising approach is the development of small molecules that catalyze the clearance of reactive aldehydes [182].

Indirect antioxidants augment the redox response without being antioxidants themselves. For example, sulforaphane, resveratrol, and curcumin induce nuclear translocation of Nrf2, a transcription factor responsible for the production of a wide array of antioxidants, and attenuate nociceptive hypersensitivity in neuropathic pain models [21,183–186]. Non-pharmacological approaches may also function in this capacity. For example, exercise increases Nrf2 expression and promotes the expression of antioxidants in the CNS as well as peripherally [187–189]. Consequently, voluntary wheel running has been shown to both prevent and reverse neuropathic pain [187,190].

Finally, ROS have a role in normal physiological processes [134–136], and there is some evidence that ROS may have protective effects after injury. For example, inflammation induced by endotoxin is exacerbated in NADPH-impaired mice relative to their wild-type counterparts [191]. In another study, yeast survival to hydrogen peroxide stress was dependent on superoxide [192]. Further work will be necessary to determine whether ROS may also have a protective role after sterile nervous system injury. However, agents have been developed to spare superoxide (e.g., peroxynitrite decomposition catalysts SRI110 and SRI6 [15]), and such approaches may prove to be important for restoring homeostasis after nervous system injury.

Concluding Remarks

Nitroxidative species are generated by mitochondria and by NOX and NOS enzymes. They enhance neuroexcitability in pain pathways through direct neuronal interactions, and indirectly by impairing mitochondria and inducing neuroinflammation. Normalizing nitroxidative signaling may be an alternative strategy to help to alleviate the enormous burden of pathological pain, which affects ~20% of the population and is poorly treated [11,193,194]. There are several

Outstanding Questions

How ubiquitous are nitroxidative signaling mechanisms within the neuraxis, beyond the classical sites already tested (peripheral nerve injury site, DRG, spinal cord)?

Are nitroxidative signaling mechanisms common or different between different preclinical pain models?

What is the relationship between the antioxidant and anti-inflammatory cytokine systems?

Do indirect antioxidants have improved translational potential for treatment of pathological pain?

areas of basic science research that may move us towards that goal (see Outstanding Questions).

Despite the extensive research implicating nitroxidative species in pathological pain states, no studies to date have quantified the crucial relationships between real-time local cellular creation of nitroxidative species, their concentration at the effect site, and the distribution of their direct effects. This challenge has not been overcome owing to the volatility of these nitroxidative species and hence their very short lifetimes *in vivo* and *ex vivo*. Several new technologies are being developed to address these issues and are discussed in **Box 1** and **Table 1**.

Lessons from the failure of direct antioxidants to improve clinical disease need to be recognized within the pain field: the effects of direct antioxidants on preclinical pain models continue to be reported despite the strong probability that the results will not translate clinically. Several studies suggest that more-robustly engaging antioxidant systems after injury can help alleviate to pain: for instance, in animal pain models, increasing the action of master antioxidant transcription factors Nrf2 or FoxO, or activating the heme-oxygenase system, show promising pain-relieving effects. Future studies could explore whether combinatorial strategies – aimed at boosting multiple antioxidants or targeting both antioxidant and nitroxidative systems simultaneously – dampen inflammation and pain. Nitroxidant dysregulation clearly contributes to neuropathology; thus, discovering new targets and therapies that restore nitroxidative balance could help to relieve pathological pain.

Box 1. New and Emerging Tools To Study Nitroxidative Species

Colorimetric and fluorescent methods for detecting the ‘shadow’ of the presence of nitroxidative species production is well established by the quantification of attendant cellular events {e.g., oxidative stress such as lipid peroxidation: thiobarbituric acid reactive substances (TBARS) [195]; DNA damage (8-oxoguanine, 8-OxoG) [196,197]} or the quantification of more stable metabolites (e.g., nitrite/nitrate using the Griess reaction [198]). These methods are not only limited in their temporal and spatial resolution, but also because of their insufficient ability to define the concentrations and time-courses of specific nitroxidative species. Establishing differential regulation of distinct nitroxidative species would be useful because specific oxygen or nitrogen species have unique outcomes in the neuroinflammatory responses. A recent example demonstrated that specifically targeting peroxynitrite reduced inflammatory progression via NLRP3 inflammasome-dependent IL-1 β /IL-18 release following intracerebral hemorrhage injury [199]. Thus, new biosensors will be necessary to improve our mechanistic understanding of how nitroxidative species affect the nervous system.

The chemistry of fluorescent probes for specific detection of both *ex vivo* and *in vivo* production of nitroxidative species has grown rapidly. A range of approaches and hence biosensors have been created that exploit platform sensing modalities, such as photoinduced electron transfer (PET) and Förster resonance energy transfer (FRET) signaling. In addition, composite biosensors that incorporate a sensor functionalized to a nanoparticle [gold particles, upconversion nanoparticles (UCNPs), and QDots] are being used to detect and/or measure ROS/RNS species (detailed in **Table 1**). Such ROS probes can quantify hypochlorite [200,201], hydroxyl [202,203], superoxide [204], hydrogen peroxide [205], and singlet oxygen [206]. Biosensors for nitric oxide [207,208], nitroxyl [209–211], peroxy nitrite [212] are also being developed.

These probes detect targeted species in cell lines, in *ex vivo* tissue, or in *in vivo* models of inflammation. However, these biosensor tools require further optimization. Further refinement of biosensors will help to improve the stability of the probe, the brightness of the fluorescing molecule, the specificity to defined species, the sensitivity of detection, and the consumption of the probe in the sensing process. However, real-time visualization and/or quantification of nitroxidative species within the CNS of a behaving preclinical rodent model of pathological pain remains an elusive goal.

The ultimate nitroxidative species biosensor would have real-time sensing capacity, with signal brightness that can detect the subcellular localization of the nitroxidative species; ideally, this probe would not be consumed/bleached in the sensing process, thereby allowing repeated measurements *in vivo*. Next-generation probes will address some of these limitations. For instance, a redox-sensitive fluorescent protein (rxRFP1), whose fluorescence intensity is positively related to the extent of oxidation of the probe, can detect varying amounts of oxidative stress within separate cellular compartments [213]. Further refining these tools will enable an improved understanding of how particular species contribute to oxidative or nitrosative stress, and will allow researchers to define how the spatiotemporal regulation of nitroxidative activity contributes to pathological pain.

Table 1. Probes Able To Detect Specific ROS and RNS Species *In Vitro*, *Ex Vivo*, or *In Vivo*^a

	Probe type	Imaging platform used	Tested <i>in vitro</i>	Tested <i>ex vivo</i> / <i>in vivo</i>	ROS/RNS stimulation	Refs
<i>ROS species</i>						
Hypochlorite	Iridium (III) complex-based two-photon phosphorescent probe	Two-photon laser scanning fluorescence microscope and confocal laser microscope	HeLa cells/RAW 264.7 cells	Zebrafish	10 mM NaClO (HeLa) 1 mg/ml LPS (RAW 264.7 and zebrafish)	[201]
Hypochlorite	Rhodamine-based hydrazide protein fluorescent probe	Fluorescence microscope	HeLa cells		50 μM OCl ⁻	[200]
Hydroxyl	Ratiometric fluorescence biosensor (gold particles conjugated with organic fluorophore)	Confocal laser microscope	HeLa cells		10 μg/ml LPS	[203]
Hydroxyl	Ratiometric fluorescence biosensor (upconversion nanoparticles conjugated with organic fluorophore)	Fluorescence microscope equipped with 980 nm laser	HeLa cells	Mouse liver	500 ng/ml PMA (<i>in vitro</i>) 1–4 mg LPS/100 g body weight (<i>in vivo</i>)	[202]
Superoxide	Fluorescein protein-based fluorescent probe	Confocal laser microscope	HCT116/BV-2/RAW 264.7 cells	Zebrafish	500 ng/ml LPS and 50 ng/ml IFN-γ (<i>in vitro</i>) PMA 200 ng/ml or antimycin A 500 nM (<i>in vivo</i>)	[204]
Hydrogen peroxide	Chemoselective fluorescent naphthylimide peroxide probe	Two-photon laser scanning fluorescence microscope	RAW 264.7 cells	Mouse lung and skin	1 μg/ml LPS (<i>in vitro</i>) 20 μg LPS (<i>in vivo</i>)	[205]
Singlet oxygen	Far-red silicon-rhodamine-based chemical fluorescent probe	Fluorescence microscope with 640 nm laser	HeLa cells/RAW 264.7 cells		Photosensitizers: 150 μg/ml 5-ALA and 5 μM TMPyP4	[206]
<i>RNS species</i>						
Nitric oxide	Chemoselective copper(II)-based fluorescence probe	Confocal laser microscope	HeLa cells/RAW 264.7 cells		50–200 μM DEA-NONOate (HeLa) 200 ng/ml LPS (RAW 264.7)	[208]
Nitric oxide	Far-red two-photon chemical fluorescent probe	Two-photon laser scanning fluorescence microscope	HeLa cells/RAW 264.7 cells	Mouse liver	25 μM NOC-9 (HeLa) 20 μg/ml LPS, 200 U/ml IFN-γ and 0.5 mg/ml L- arginine (RAW 264.7) 1–4 mg/ml LPS (<i>in vivo</i>)	[207]
Nitroxyl	FRET-based ratiometric chemical fluorescent probe	Confocal laser microscope	HeLa cells		100 μM AS	[211]

Table 1. (continued)

	Probe type	Imaging platform used	Tested <i>in vitro</i>	Tested <i>ex vivo</i> / <i>in vivo</i>	ROS/RNS stimulation	Refs
Nitroxyl	Near infrared chemical fluorescent probe	Confocal laser microscope and <i>in vivo</i> imaging system	RAW 264.7 cells	Mouse (<i>in vivo</i>)	100 µM AS (RAW 264.7) 500 µM AS (i.p. Mouse)	[210]
Nitroxyl	Lysosome-targetable near infrared chemical fluorescent probe	Confocal laser microscope and <i>in vivo</i> imaging system	RAW 264.7 cells	Mouse (<i>in vivo</i>)	200 µM AS (RAW 264.7) 1 mM AS (i.p. mouse)	[209]
Peroxynitrite	Boronate-based chemical fluorescent probe	Confocal laser microscope	HeLa cells/RAW 264.7 cells		5 and 20 µM peroxynitrite solution (HeLa) 1 µg/ml LPS, 50 ng/ml IFN-γ, 2.5 ng/ml PMA (RAW 264.7)	[212]

^aAbbreviations: AS, Angel's salt (nitroxyl donor); 5-ALA, 5-aminolevulinic acid (drug used in photodynamic therapy, known to produce singlet oxygen); antimycin A, produces endogenous ROS/RNS by driving apoptosis; DEA-NOONOate, 2-(*N,N*-diethylamino)-diazenol 2-oxide (nitric oxide donor); IFN-γ, interferon γ (produces endogenous ROS/RNS); LPS, lipopolysaccharide (produces endogenous ROS/RNS); NOC-9, 6-(2-hydroxy-1-methyl-2-nitrosohydrazino)-*N*-methyl-1-hexanamine (nitric oxide donor); PMA, phorbol 12-myristate-13-acetate (activates protein kinase C *in vivo* and *in vitro*); TMPyP4, 5,10,15,20-tetra-(*N*-methyl-4-pyridyl)porphyrin (drug used in photodynamic therapy, known to produce singlet oxygen).

Acknowledgments

This work was supported by an Australian Postgraduate Research scholarship (V.S.), an ARC Research Fellowship (DP110100297; M.R.H.), funding from the ARC Centre of Excellence for Nanoscale Biophotonics (CE140100003; M.R.H.), National Institutes of Health (NIH) grant DE021966 (L.R.W.), and a Wings for Life Project Grant (L.R.W./A.D.G.).

References

- Krebs, H.A. and Johnson, W.A. (1937) The role of citric acid in intermediate metabolism in animal tissues. *Enzymologia* 4, 148–156
- Gozsy, B. and Szent-Gyorgyi, A. (1934) On the mechanism of primary respiration in pigeon breast muscle. *Hoppe-Seyler's Z. Physiol. Chem.* 224, 1–10 (in German)
- Harden, A. and Young, W.J. (1906) The alcoholic ferment of yeast-juice. *Proc. R. Soc. Lond. B Biol. Sci.* 77, 405–420
- Maass, O. and Hatcher, W.H. (1920) The properties of pure hydrogen peroxide. I. *J Am Chem Soc* 42, 2548–2569
- Grace, P.M. et al. (2014) Pathological pain and the neuroimmune interface. *Nat. Rev. Immunol.* 14, 217–231
- Salvermini, D. et al. (2011) Roles of reactive oxygen and nitrogen species in pain. *Free Radic. Biol. Med.* 51, 951–966
- Janes, K. et al. (2012) Anti-superoxide and anti-peroxynitrite strategies in pain suppression. *Biochim. Biophys. Acta* 1822, 815–821
- Little, J.W. et al. (2012) Reactive nitroxidative species and nociceptive processing: determining the roles for nitric oxide, superoxide, and peroxynitrite in pain. *Amino Acids* 42, 75–94
- Symons-Liguori, A. et al. (2016) The contribution of nitroxidative stress to pathophysiological pain and opioid analgesic failure. In *Redox-Active Therapeutics* (Batinic-Haberle, I. et al., eds), pp. 563–595, Springer International Publishing
- Batinic-Haberle, I. et al. (2016) Mn porphyrin-based redox-active therapeutics. In *Redox-Active Therapeutics* (Batinic-Haberle, I. et al., eds), pp. 165–211, Springer International Publishing
- Pizzo, P.A. and Clark, N.M. (2012) Alleviating suffering 101 – pain relief in the United States. *N. Engl. J. Med.* 366, 197–199
- Burma, N.E. et al. (2016) Animal models of chronic pain: advances and challenges for clinical translation. *J. Neurosci. Res.* Published online July 4, 2016. <http://dx.doi.org/10.1002/jnr.23768>
- Casas, A.I. et al. (2015) Reactive oxygen-related diseases: therapeutic targets and emerging clinical indications. *Antioxid. Redox Signal.* 23, 1171–1185
- Holmström, K.M. and Finkel, T. (2014) Cellular mechanisms and physiological consequences of redox-dependent signalling. *Nat. Rev. Mol. Cell Biol.* 15, 411–421
- Doyle, T. et al. (2012) Targeting the overproduction of peroxynitrite for the prevention and reversal of paclitaxel-induced neuropathic pain. *J. Neurosci.* 32, 6149–6160
- Hassler, S.N. et al. (2014) Reactive oxygen species and lipid peroxidation inhibitors reduce mechanical sensitivity in a chronic neuropathic pain model of spinal cord injury in rats. *J. Neurochem.* 131, 413–417
- Nayernia, Z. et al. (2014) New insights on NOX enzymes in the central nervous system. *Antioxid. Redox Signal.* 20, 2815–2837
- Ibi, M. et al. (2008) Reactive oxygen species derived from NOX1/NADPH oxidase enhance inflammatory pain. *J. Neurosci.* 28, 9486–9494
- Kim, J.-S. et al. (2010) Glycogen synthase kinase 3beta and beta-catenin pathway is involved in Toll-like receptor 4-mediated NADPH oxidase 1 expression in macrophages. *FEBS J.* 277, 2830–2837
- Kallenborn-Gerhardt, W. et al. (2014) Nox2-dependent signaling between macrophages and sensory neurons contributes to neuropathic pain hypersensitivity. *Pain* 155, 2161–2170
- Kim, D. et al. (2010) NADPH oxidase 2-derived reactive oxygen species in spinal cord microglia contribute to peripheral nerve

- injury-induced neuropathic pain. *Proc. Natl. Acad. Sci. U.S.A.* 107, 14851–14856
22. Hackel, D. *et al.* (2013) The connection of monocytes and reactive oxygen species in pain. *PLoS ONE* 8, e63564
 23. Kallenborn-Gerhardt, W. *et al.* (2012) NADPH oxidase-4 maintains neuropathic pain after peripheral nerve injury. *J. Neurosci.* 32, 10136–10145
 24. Lee, C.F. *et al.* (2010) Nox4 is a novel inducible source of reactive oxygen species in monocytes and macrophages and mediates oxidized low density lipoprotein-induced macrophage death. *Circ. Res.* 106, 1489–1497
 25. Grace, P.M. *et al.* (2010) A novel animal model of graded neuropathic pain: utility to investigate mechanisms of population heterogeneity. *J. Neurosci. Methods* 193, 47–53
 26. Wen, Y-R. *et al.* (2007) Nerve conduction blockade in the sciatic nerve prevents but does not reverse the activation of p38 mitogen-activated protein kinase in spinal microglia in the rat spared nerve injury model. *Anesthesiology* 107, 312–321
 27. Xie, W. *et al.* (2009) Early blockade of injured primary sensory afferents reduces glial cell activation in two rat neuropathic pain models. *Neuroscience* 160, 847–857
 28. Cury, Y. *et al.* (2011) Pain and analgesia: the dual effect of nitric oxide in the nociceptive system. *Nitric Oxide Biol. Chem.* 25, 243–254
 29. Bredt, D.S. *et al.* (1990) Localization of nitric oxide synthase indicating a neural role for nitric oxide. *Nature* 347, 768–770
 30. Cao, J. *et al.* (2005) The PSD95-nNOS interface: a target for inhibition of excitotoxic p38 stress-activated protein kinase activation and cell death. *J. Cell Biol.* 168, 117–126
 31. Askwith, T. *et al.* (2012) Taurine reduces nitrosative stress and nitric oxide synthase expression in high glucose-exposed human Schwann cells. *Exp. Neurol.* 233, 154–162
 32. Annedi, S.C. *et al.* (2011) Discovery of N-(3-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)-1H-indol-6-yl) thiophene-2-carboximide as a selective inhibitor of human neuronal nitric oxide synthase (nNOS) for the treatment of pain. *J. Med. Chem.* 54, 7408–7416
 33. Keilhoff, G. *et al.* (2013) Time-course of neuropathic pain in mice deficient in neuronal or inducible nitric oxide synthase. *Neurosci. Res.* 77, 215–221
 34. Mihara, Y. *et al.* (2011) Involvement of spinal NR2B-containing NMDA receptors in oxaliplatin-induced mechanical allodynia in rats. *Mol. Pain* 7, 8
 35. Mukherjee, P. *et al.* (2014) Development of nitric oxide synthase inhibitors for neurodegeneration and neuropathic pain. *Chem. Soc. Rev.* 43, 6814–6838
 36. Bonnefous, C. *et al.* (2009) Discovery of inducible nitric oxide synthase (iNOS) inhibitor development candidate KD7332. Part 1. Identification of a novel, potent, and selective series of quinolinone iNOS dimerization inhibitors that are orally active in rodent pain models. *J. Med. Chem.* 52, 3047–3062
 37. Tanabe, M. *et al.* (2009) Pharmacological assessments of nitric oxide synthase isoforms and downstream diversity of NO signaling in the maintenance of thermal and mechanical hypersensitivity after peripheral nerve injury in mice. *Neuropharmacology* 56, 702–708
 38. Borsig, E. *et al.* (2013) Endothelial nitric oxide synthase in dorsal root ganglia during chronic inflammatory nociception. *Cells Tissues Organs* 197, 159–168
 39. Galluzzi, L. *et al.* (2012) Mitochondria: master regulators of danger signalling. *Nat. Rev. Mol. Cell Biol.* 13, 780–788
 40. Schwartz, E.S. *et al.* (2008) Oxidative stress in the spinal cord is an important contributor in capsaicin-induced mechanical secondary hyperalgesia in mice. *Pain* 138, 514–524
 41. Schwartz, E.S. *et al.* (2009) Persistent pain is dependent on spinal mitochondrial antioxidant levels. *J. Neurosci.* 29, 159–168
 42. Chu, C. *et al.* (2011) Mitochondrial dependence of nerve growth factor-induced mechanical hyperalgesia. *Pain* 152, 1832–1837
 43. Ferrari, L.F. and Levine, J.D. (2010) Alcohol consumption enhances antiretroviral painful peripheral neuropathy by mitochondrial mechanisms. *Eur. J. Neurosci.* 32, 811–818
 44. Joseph, E.K. and Levine, J.D. (2006) Mitochondrial electron transport in models of neuropathic and inflammatory pain. *Pain* 121, 105–114
 45. Joseph, E.K. and Levine, J.D. (2010) Multiple PKC ϵ -dependent mechanisms mediating mechanical hyperalgesia. *Pain* 150, 17–21
 46. Zheng, H. *et al.* (2011) Functional deficits in peripheral nerve mitochondria in rats with paclitaxel- and oxaliplatin-evoked painful peripheral neuropathy. *Exp. Neurol.* 232, 154–161
 47. Basbaum, A.I. *et al.* (2009) Cellular and molecular mechanisms of pain. *Cell* 139, 267–284
 48. Gold, M.S. and Gebhart, G.F. (2010) Nociceptor sensitization in pain pathogenesis. *Nat. Med.* 16, 1248–1257
 49. Latremoliere, A. and Woolf, C.J. (2009) Central sensitization: a generator of pain hypersensitivity by central neural plasticity. *J. Pain* 10, 895–926
 50. Ren, K. and Dubner, R. (2010) Interactions between the immune and nervous systems in pain. *Nat. Med.* 16, 1267–1276
 51. Gwak, Y.S. *et al.* (2013) Reactive oxygen species contribute to neuropathic pain and locomotor dysfunction via activation of CaMKII in remote segments following spinal cord contusion injury in rats. *Pain* 154, 1699–1708
 52. Ndengel, M.M. *et al.* (2008) Cyclooxygenases 1 and 2 contribute to peroxynitrite-mediated inflammatory pain hypersensitivity. *FASEB J.* 22, 3154–3164
 53. Yowtak, J. *et al.* (2011) Reactive oxygen species contribute to neuropathic pain by reducing spinal GABA release. *Pain* 152, 844–852
 54. Wang, Z-Q. *et al.* (2004) A newly identified role for superoxide in inflammatory pain. *J. Pharmacol. Exp. Ther.* 309, 869–878
 55. Sözbir, E. and Naziroğlu, M. (2016) Diabetes enhances oxidative stress-induced TRPM2 channel activity and its control by N-acetylcysteine in rat dorsal root ganglion and brain. *Metab. Brain Dis.* 31, 385–393
 56. Chen, Z. *et al.* (2010) NMDA-receptor activation and nitrooxidative regulation of the glutamatergic pathway during nociceptive processing. *Pain* 149, 100–106
 57. Lee, I. *et al.* (2007) The role of reactive oxygen species in capsaicin-induced mechanical hyperalgesia and in the activities of dorsal horn neurons. *Pain* 133, 9–17
 58. Rausaria, S. *et al.* (2011) Retooling manganese(III) porphyrin-based peroxynitrite decomposition catalysts for selectivity and oral activity: a potential new strategy for treating chronic pain. *J. Med. Chem.* 54, 8658–8669
 59. Lorenz, J.E. *et al.* (2014) Oxidant-induced activation of cGMP-dependent protein kinase I α mediates neuropathic pain after peripheral nerve injury. *Antioxid. Redox Signal.* 21, 1504–1515
 60. Luo, C. *et al.* (2012) Presynaptically localized cyclic GMP-dependent protein kinase 1 is a key determinant of spinal synaptic potentiation and pain hypersensitivity. *PLoS Biol.* 10, e1001283
 61. Muscoli, C. *et al.* (2013) Posttranslational nitration of tyrosine residues modulates glutamate transmission and contributes to N-methyl-D-aspartate-mediated thermal hyperalgesia. *Mediators Inflamm.* 2013, 950947
 62. Gao, X. *et al.* (2007) Reactive oxygen species (ROS) are involved in enhancement of NMDA-receptor phosphorylation in animal models of pain. *Pain* 131, 262–271
 63. Zanelli, S.A. *et al.* (2002) Nitration is a mechanism of regulation of the NMDA receptor function during hypoxia. *Neuroscience* 112, 869–877
 64. Yowtak, J. *et al.* (2013) Effect of antioxidant treatment on spinal GABA neurons in a neuropathic pain model in the mouse. *Pain* 154, 2469–2476
 65. Julius, D. (2013) TRP channels and pain. *Annu. Rev. Cell Dev. Biol.* 29, 355–384
 66. Xanthos, D.N. and Sandkühler, J. (2014) Neurogenic neuroinflammation: inflammatory CNS reactions in response to neuronal activity. *Nat. Rev. Neurosci.* 15, 43–53
 67. Andersson, D.A. *et al.* (2015) Streptozotocin stimulates the ion channel TRPA1 directly: involvement of peroxynitrite. *J. Biol. Chem.* 290, 15185–15196

68. Bautista, D.M. *et al.* (2006) TRPA1 mediates the inflammatory actions of environmental irritants and proalgesic agents. *Cell* 124, 1269–1282
69. Bessac, B.F. *et al.* (2009) Transient receptor potential ankyrin 1 antagonists block the noxious effects of toxic industrial isocyanates and tear gases. *FASEB J.* 23, 1102–1114
70. Taylor-Clark, T.E. and Undem, B.J. (2010) Ozone activates airway nerves via the selective stimulation of TRPA1 ion channels. *J. Physiol.* 588, 423–433
71. Taylor-Clark, T.E. *et al.* (2009) Nitrooleic acid, an endogenous product of nitrative stress, activates nociceptive sensory nerves via the direct activation of TRPA1. *Mol. Pharmacol.* 75, 820–829
72. Trevisani, M. *et al.* (2007) 4-Hydroxyonenal, an endogenous aldehyde, causes pain and neurogenic inflammation through activation of the irritant receptor TRPA1. *Proc. Natl. Acad. Sci.* 104, 13519–13524
73. Due, M.R. *et al.* (2014) Acrolein involvement in sensory and behavioral hypersensitivity following spinal cord injury in the rat. *J. Neurochem.* 128, 776–786
74. Chen, Z. *et al.* (2016) Mitigation of sensory and motor deficits by acrolein scavenger phenelzine in a rat model of spinal cord contusive injury. *J. Neurochem.* 138, 328–338
75. Trevisani, G. *et al.* (2013) Novel therapeutic strategy to prevent chemotherapy-induced persistent sensory neuropathy by TRPA1 blockade. *Cancer Res.* 73, 3120–3131
76. Chen, S. *et al.* (2013) Role of TRPM2 in cell proliferation and susceptibility to oxidative stress. *Am. J. Physiol. Cell Physiol.* 304, C548–C560
77. Haraguchi, K. *et al.* (2012) TRPM2 contributes to inflammatory and neuropathic pain through the aggravation of pronociceptive inflammatory responses in mice. *J. Neurosci.* 32, 3931–3941
78. Naziroğlu, M. *et al.* (2013) Role of TRPM2 cation channels in dorsal root ganglion of rats after experimental spinal cord injury. *Muscle Nerve* 48, 945–950
79. Özdemir, Ü.S. *et al.* (2015) *Hypericum perforatum* attenuates spinal cord injury-induced oxidative stress and apoptosis in the dorsal root ganglion of rats: involvement of TRPM2 and TRPV1 channels. *Mol. Neurobiol.* 53, 3540–3551
80. Perraud, A-L. *et al.* (2005) Accumulation of free ADP-ribose from mitochondria mediates oxidative stress-induced gating of TRPM2 cation channels. *J. Biol. Chem.* 280, 6138–6148
81. Yamamoto, S. *et al.* (2008) TRPM2-mediated Ca^{2+} influx induces chemokine production in monocytes that aggravates inflammatory neutrophil infiltration. *Nat. Med.* 14, 738–747
82. Isami, K. *et al.* (2013) Involvement of TRPM2 in peripheral nerve injury-induced infiltration of peripheral immune cells into the spinal cord in mouse neuropathic pain model. *PLOS ONE* 8, e66410
83. Chung, M-K. *et al.* (2015) The role of TRPM2 in hydrogen peroxide-induced expression of inflammatory cytokine and chemokine in rat trigeminal ganglia. *Neuroscience* 297, 160–169
84. Naziroğlu, M. *et al.* (2014) Modulation of oxidative stress and Ca^{2+} mobilization through TRPM2 channels in rat dorsal root ganglion neuron by *Hypericum perforatum*. *Neuroscience* 263, 27–35
85. So, K. *et al.* (2015) Involvement of TRPM2 in a wide range of inflammatory and neuropathic pain mouse models. *J. Pharmacol. Sci.* 127, 237–243
86. Chuang, H. and Lin, S. (2009) Oxidative challenges sensitize the capsaicin receptor by covalent cysteine modification. *Proc. Natl. Acad. Sci. U.S.A.* 106, 20097–20102
87. Susankova, K. *et al.* (2006) Reducing and oxidizing agents sensitize heat-activated vanilloid receptor (TRPV1) current. *Mol. Pharmacol.* 70, 383–394
88. Westlund, K.N. *et al.* (2010) Impact of central and peripheral TRPV1 and ROS levels on proinflammatory mediators and nociceptive behavior. *Mol. Pain* 6, 46
89. Patwardhan, A.M. *et al.* (2009) Activation of TRPV1 in the spinal cord by oxidized linoleic acid metabolites contributes to inflammatory hyperalgesia. *Proc. Natl. Acad. Sci. U.S.A.* 106, 18820–18824
90. Patwardhan, A.M. *et al.* (2010) Heat generates oxidized linoleic acid metabolites that activate TRPV1 and produce pain in rodents. *J. Clin. Invest.* 120, 1617–1626
91. Bennett, G.J. *et al.* (2014) Mitotoxicity in distal symmetrical sensory peripheral neuropathies. *Nat. Rev. Neurol.* 10, 326–336
92. MacMillan-Crow, L.A. and Thompson, J.A. (1999) Tyrosine modifications and inactivation of active site manganese superoxide dismutase mutant (Y34F) by peroxynitrite. *Arch. Biochem. Biophys.* 366, 82–88
93. Zarkovic, N. *et al.* (2013) Pathophysiological relevance of aldehydic protein modifications. *J. Proteomics* 92, 239–247
94. Sinha, K. *et al.* (2013) Oxidative stress: the mitochondria-dependent and mitochondria-independent pathways of apoptosis. *Arch. Toxicol.* 87, 1157–1180
95. Yuan, H. *et al.* (2006) Mitochondrial fission is an upstream and required event for bax foci formation in response to nitric oxide in cortical neurons. *Cell Death Differ.* 14, 462–471
96. Scheving, R. *et al.* (2012) Protein S-nitrosylation and denitrosylation in the mouse spinal cord upon injury of the sciatic nerve. *J. Proteomics* 75, 3987–4004
97. Joseph, E.K. and Levine, J.D. (2004) Caspase signalling in neuropathic and inflammatory pain in the rat. *Eur. J. Neurosci.* 20, 2896–2902
98. Ferrari, L.F. *et al.* (2011) Role of Drp1, a key mitochondrial fission protein, in neuropathic pain. *J. Neurosci.* 31, 11404–11410
99. Janes, K. *et al.* (2013) Bioenergetic deficits in peripheral nerve sensory axons during chemotherapy-induced neuropathic pain resulting from peroxynitrite-mediated post-translational nitration of mitochondrial superoxide dismutase. *Pain* 154, 2432–2440
100. Lim, T.K.Y. *et al.* (2015) Mitochondrial and bioenergetic dysfunction in trauma-induced painful peripheral neuropathy. *Mol. Pain* 11, 58
101. Austin, P.J. and Moalem-Taylor, G. (2010) The neuro-immune balance in neuropathic pain: involvement of inflammatory immune cells, immune-like glial cells and cytokines. *J. Neuroimmunol.* 229, 26–50
102. Grace, P.M. *et al.* (2011) Peripheral immune contributions to the maintenance of central glial activation underlying neuropathic pain. *Brain. Behav. Immun.* 25, 1322–1332
103. Gaudet, A.D. *et al.* (2011) Wallerian degeneration: gaining perspective on inflammatory events after peripheral nerve injury. *J. Neuroinflammation* 8, 110
104. Genovese, T. *et al.* (2009) Effects of a metalloporphyrinic peroxynitrite decomposition catalyst, ww-85, in a mouse model of spinal cord injury. *Free Radic. Res.* 43, 631–645
105. Kamata, H. *et al.* (2005) Reactive oxygen species promote TNFalpha-induced death and sustained JNK activation by inhibiting MAP kinase phosphatases. *Cell* 120, 649–661
106. Matata, B.M. and Galihanes, M. (2002) Peroxynitrite is an essential component of cytokines production mechanism in human monocytes through modulation of nuclear factor-kappa B DNA binding activity. *J. Biol. Chem.* 277, 2330–2335
107. Zhou, H. *et al.* (2013) CD11b/CD18 (Mac-1) is a novel surface receptor for extracellular double-stranded RNA to mediate cellular inflammatory responses. *J. Immunol.* 190, 115–125
108. Li, J. *et al.* (2016) N-acetyl-cysteine attenuates neuropathic pain by suppressing matrix metalloproteinases. *Pain* 157, 1711–1723
109. Zhang, Q. *et al.* (2010) Circulating mitochondrial DAMPs cause inflammatory responses to injury. *Nature* 464, 104–107
110. Lee, J.H. *et al.* (2013) Interaction of NADPH oxidase 1 with Toll-like receptor 2 induces migration of smooth muscle cells. *Cardiovasc. Res.* 99, 483–493
111. Nakahira, K. *et al.* (2006) Carbon monoxide differentially inhibits TLR signaling pathways by regulating ROS-induced trafficking of TLRs to lipid rafts. *J. Exp. Med.* 203, 2377–2389
112. Park, H.S. *et al.* (2004) Direct interaction of TLR4 with NAD(P)H oxidase 4 isozyme is essential for lipopolysaccharide-induced production of reactive oxygen species and activation of NF- κ B. *J. Immunol.* 173, 3589–3595
113. Yang, C-S. *et al.* (2008) ASK1-p38 MAPK-p47phox activation is essential for inflammatory responses during tuberculosis via TLR2-ROS signalling. *Cell. Microbiol.* 10, 741–754

114. Yang, C-S. *et al.* (2009) NADPH oxidase 2 interaction with TLR2 is required for efficient innate immune responses to mycobacteria via cathelicidin expression. *J. Immunol.* 182, 3696–3705
115. Matsuzawa, A. *et al.* (2005) ROS-dependent activation of the TRAF6-ASK1-p38 pathway is selectively required for TLR4-mediated innate immunity. *Nat. Immunol.* 6, 587–592
116. Wong, S.W. *et al.* (2009) Fatty acids modulate Toll-like receptor 4 activation through regulation of receptor dimerization and recruitment into lipid rafts in a reactive oxygen species-dependent manner. *J. Biol. Chem.* 284, 27384–27392
117. Lochhead, J.J. *et al.* (2012) Tempol modulates changes in xenobiotic permeability and occludin oligomeric assemblies at the blood–brain barrier during inflammatory pain. *Am. J. Physiol. Heart Circ. Physiol.* 302, H582–593
118. Zhou, R. *et al.* (2010) Thioredoxin-interacting protein links oxidative stress to inflammasome activation. *Nat. Immunol.* 11, 136–140
119. Latz, E. *et al.* (2013) Activation and regulation of the inflammasomes. *Nat. Rev. Immunol.* 13, 397–411
120. Grace, P.M. *et al.* (2016) Morphine paradoxically prolongs neuropathic pain in rats by amplifying spinal NLRP3 inflammasome activation. *Proc. Natl. Acad. Sci. U.S.A.* 113, E3441–E3450
121. Nakahira, K. *et al.* (2011) Autophagy proteins regulate innate immune responses by inhibiting the release of mitochondrial DNA mediated by the NALP3 inflammasome. *Nat. Immunol.* 12, 222–230
122. Shimada, K. *et al.* (2012) Oxidized mitochondrial DNA activates the NLRP3 inflammasome during apoptosis. *Immunity* 36, 401–414
123. Zhou, R. *et al.* (2011) A role for mitochondria in NLRP3 inflammasome activation. *Nature* 469, 221–225
124. Zhong, Z. *et al.* (2013) TRPM2 links oxidative stress to NLRP3 inflammasome activation. *Nat. Commun.* 4, 1611
125. Anrather, J. *et al.* (2006) NF-kappaB regulates phagocytic NADPH oxidase by inducing the expression of gp91phox. *J. Biol. Chem.* 281, 5657–5667
126. Guo, Z. *et al.* (2007) Identification of a classic cytokine-induced enhancer upstream in the human iNOS promoter. *FASEB J.* 21, 535–542
127. Lee, J-G. *et al.* (2008) A combination of Lox-1 and Nox1 regulates TLR9-mediated foam cell formation. *Cell. Signal.* 20, 2266–2275
128. Li, Y. *et al.* (2007) Regulation of neuronal nitric oxide synthase exon 1f gene expression by nuclear factor-kappaB acetylation in human neuroblastoma cells. *J. Neurochem.* 101, 1194–1204
129. Yoo, B.K. *et al.* (2008) Activation of p38 MAPK induced peroxynitrite generation in LPS plus IFN-gamma-stimulated rat primary astrocytes via activation of iNOS and NADPH oxidase. *Neurochem. Int.* 52, 1188–1197
130. Sorge, R.E. *et al.* (2012) Genetically determined P2X7 receptor pore formation regulates variability in chronic pain sensitivity. *Nat. Med.* 18, 595–599
131. Apolloni, S. *et al.* (2013) The NADPH oxidase pathway is dysregulated by the P2X7 receptor in the SOD1-G93A microglia model of amyotrophic lateral sclerosis. *J. Immunol.* 190, 5187–5195
132. Kim, S.Y. *et al.* (2007) ATP released from beta-amyloid-stimulated microglia induces reactive oxygen species production in an autocrine fashion. *Exp. Mol. Med.* 39, 820–827
133. Parvathenani, L.K. *et al.* (2003) P2X7 mediates superoxide production in primary microglia and is up-regulated in a transgenic mouse model of Alzheimer's disease. *J. Biol. Chem.* 278, 13309–13317
134. Sena, L.A. and Chandel, N.S. (2012) Physiological roles of mitochondrial reactive oxygen species. *Mol. Cell* 48, 158–167
135. Zuo, L. *et al.* (2015) Biological and physiological role of reactive oxygen species – the good, the bad and the ugly. *Acta Physiol.* 214, 329–348
136. Massaad, C.A. and Klann, E. (2011) Reactive reactive oxygen species in the regulation of synaptic plasticity and memory. *Antioxid. Redox Signal.* 14, 2013–2054
137. Dang, J. *et al.* (2012) Nrf2 expression by neurons, astroglia, and microglia in the cerebral cortical penumbra of ischemic rats. *J. Mol. Neurosci.* 46, 578–584
138. Ishii, T. *et al.* (1999) Oxidative stress-inducible proteins in macrophages. *Free Radic. Res.* 31, 351–355
139. Vincent, A.M. *et al.* (2009) Sensory neurons and Schwann cells respond to oxidative stress by increasing antioxidant defense mechanisms. *Antioxid. Redox Signal.* 11, 425–438
140. Hayes, J.D. and Dinkova-Kostova, A.T. (2014) The Nrf2 regulatory network provides an interface between redox and intermediary metabolism. *Trends Biochem. Sci.* 39, 199–218
141. Klotz, L-O. *et al.* (2015) Redox regulation of FoxO transcription factors. *Redox Biol.* 6, 51–72
142. Vilhardt, F. *et al.* (2016) Microglia antioxidant systems and redox signaling. *Br. J. Pharmacol.* Published online January 11, 2016. <http://dx.doi.org/10.1111/bph.13426>
143. Goecks, C.S.B. *et al.* (2012) Assessment of oxidative parameters in rat spinal cord after chronic constriction of the sciatic nerve. *Neurochem. Res.* 37, 1952–1958
144. Guedes, R.P. *et al.* (2006) Neuropathic pain modifies antioxidant activity in rat spinal cord. *Neurochem. Res.* 31, 603–609
145. Guedes, R.P. *et al.* (2009) Sciatic nerve transection increases glutathione antioxidant system activity and neuronal nitric oxide synthase expression in the spinal cord. *Brain Res. Bull.* 80, 422–427
146. Hu, P. *et al.* (2015) Secoisolariciresinol diglycoside, a flaxseed lignan, exerts analgesic effects in a mouse model of type 1 diabetes: engagement of antioxidant mechanism. *Eur. J. Pharmacol.* 767, 183–192
147. Kallenborn-Gerhardt, W. *et al.* (2013) Antioxidant activity of seselin 2 controls neuropathic pain after peripheral nerve injury. *Antioxid. Redox Signal.* 19, 2013–2023
148. Liu, X. *et al.* (2015) Spinal heme oxygenase-1 (HO-1) exerts antinociceptive effects against neuropathic pain in a mouse model of L5 spinal nerve ligation. *Pain Med.* 17, 220–229
149. Naik, A.K. *et al.* (2006) Role of oxidative stress in pathophysiology of peripheral neuropathy and modulation by N-acetyl-L-cysteine in rats. *Eur. J. Pain* 10, 573–579
150. Pabreja, K. *et al.* (2011) Minocycline attenuates the development of diabetic neuropathic pain: possible anti-inflammatory and anti-oxidant mechanisms. *Eur. J. Pharmacol.* 661, 15–21
151. Pathak, N.N. *et al.* (2014) Atorvastatin attenuates neuropathic pain in rat neuropathy model by down-regulating oxidative damage at peripheral, spinal and supraspinal levels. *Neurochem. Int.* 68, 1–9
152. Scheid, T. *et al.* (2013) Sciatic nerve transection modulates oxidative parameters in spinal and supraspinal regions. *Neurochem. Res.* 38, 935–942
153. Berger, J.V. *et al.* (2011) Enhanced neuroinflammation and pain hypersensitivity after peripheral nerve injury in rats expressing mutated superoxide dismutase 1. *J. Neuroinflammation* 8, 33
154. Kong, Q. *et al.* (2012) Heme oxygenase-1 inhibits neuropathic pain in rats with diabetic mellitus. *Neural Regen. Res.* 7, 2305–2311
155. Shen, Y. *et al.* (2015) Exogenous induction of HO-1 alleviates vincristine-induced neuropathic pain by reducing spinal glial activation in mice. *Neurobiol. Dis.* 79, 100–110
156. Chen, G. *et al.* (2015) Intrathecal bone marrow stromal cells inhibit neuropathic pain via TGF- β secretion. *J. Clin. Invest.* 125, 3226–3240
157. Kwiłas, A.J. *et al.* (2015) The therapeutic potential of interleukin-10 in neuroimmune diseases. *Neuropharmacology* 96, 55–69
158. Lee, T-S. and Chau, L-Y. (2002) Heme oxygenase-1 mediates the anti-inflammatory effect of interleukin-10 in mice. *Nat. Med.* 8, 240–246
159. Qian, L. *et al.* (2006) Interleukin-10 protects lipopolysaccharide-induced neurotoxicity in primary midbrain cultures by inhibiting the function of NADPH oxidase. *J. Pharmacol. Exp. Ther.* 319, 44–52
160. Qian, L. *et al.* (2008) Potent anti-inflammatory and neuroprotective effects of TGF-beta1 are mediated through the inhibition of

- ERK and p47phox-Ser345 phosphorylation and translocation in microglia. *J. Immunol.* 181, 660–668
161. Jadhav, A. et al. (2013) The heme oxygenase system selectively enhances the anti-inflammatory macrophage-M2 phenotype, reduces pericardial adiposity, and ameliorated cardiac injury in diabetic cardiomyopathy in Zucker diabetic fatty rats. *J. Pharmacol. Exp. Ther.* 345, 239–249
162. Piantadosi, C.A. et al. (2011) Heme oxygenase-1 couples activation of mitochondrial biogenesis to anti-inflammatory cytokine expression. *J. Biol. Chem.* 286, 16374–16385
163. Janes, K. et al. (2014) A3 adenosine receptor agonist prevents the development of paclitaxel-induced neuropathic pain by modulating spinal glial-restricted redox-dependent signaling pathways. *Pain* 155, 2560–2567
164. Little, J.W. et al. (2015) Endogenous adenosine A3 receptor activation selectively alleviates persistent pain states. *Brain* 138, 28–35
165. Loram, L.C. et al. (2009) Enduring reversal of neuropathic pain by a single intrathecal injection of adenosine 2A receptor agonists: a novel therapy for neuropathic pain. *J. Neurosci.* 29, 14015–14025
166. Sun, W. et al. (2007) Pharmacologic characterization of novel adenosine A2A receptor agonists in equine neutrophils. *Am. J. Vet. Res.* 68, 981–987
167. Sun, W. et al. (2007) Effects of stimulation of adenosine A2A receptors on lipopolysaccharide-induced production of reactive oxygen species by equine neutrophils. *Am. J. Vet. Res.* 68, 649–656
168. Grace, P.M. et al. (2015) Opioid-induced central immune signaling: implications for opioid analgesia. *Headache* 55, 475–489
169. Hutchinson, M.R. et al. (2011) Exploring the neuroimmunopharmacology of opioids: an integrative review of mechanisms of central immune signaling and their implications for opioid analgesia. *Pharmacol. Rev.* 63, 772–810
170. Salvermini, D. and Neumann, W.L. (2009) Peroxynitrite: a strategic linchpin of opioid analgesic tolerance. *Trends Pharmacol. Sci.* 30, 194–202
171. Doyle, T. et al. (2010) Spinal NADPH oxidase is a source of superoxide in the development of morphine-induced hyperalgesia and antinociceptive tolerance. *Neurosci. Lett.* 483, 85–89
172. Doyle, T. et al. (2013) NADPH-oxidase 2 activation promotes opioid-induced antinociceptive tolerance in mice. *Neuroscience* 241, 1–9
173. Ibi, M. et al. (2011) Involvement of NOX1/NADPH oxidase in morphine-induced analgesia and tolerance. *J. Neurosci.* 31, 18094–18103
174. Little, J.W. et al. (2013) Spinal mitochondrial-derived peroxynitrite enhances neuroimmune activation during morphine hyperalgesia and antinociceptive tolerance. *Pain* 154, 978–986
175. Muscoli, C. et al. (2007) Therapeutic manipulation of peroxynitrite attenuates the development of opiate-induced antinociceptive tolerance in mice. *J. Clin. Invest.* 117, 3530–3539
176. Batinic-Haberle, I. et al. (2009) Lipophilicity is a critical parameter that dominates the efficacy of metalloporphyrins in blocking the development of morphine antinociceptive tolerance through peroxynitrite-mediated pathways. *Free Radic. Biol. Med.* 46, 212–219
177. Little, J.W. et al. (2012) Supraspinal peroxynitrite modulates pain signaling by suppressing the endogenous opioid pathway. *J. Neurosci.* 32, 10797–10808
178. Schmidt, H.H.W. et al. (2015) Antioxidants in translational medicine. *Antioxid. Redox Signal.* 23, 1130–1143
179. Salvermini, D. et al. (2002) SOD mimetics are coming of age. *Nat. Rev. Drug Discov.* 1, 367–374
180. Slosky, L.M. et al. (2016) The cystine/glutamate antiporter system xc– drives breast tumor cell glutamate release and cancer-induced bone pain. *Pain* 157, 2605–2616
181. Dao, V.T.-V. et al. (2015) Pharmacology and clinical drug candidates in redox medicine. *Antioxid. Redox Signal.* 23, 1113–1129
182. Zambelli, V.O. et al. (2014) Aldehyde dehydrogenase-2 regulates nociception in rodent models of acute inflammatory pain. *Sci. Transl. Med.* 6, 251ra118
183. Fattori, V. et al. (2015) Curcumin inhibits superoxide anion-induced pain-like behavior and leukocyte recruitment by increasing Nrf2 expression and reducing NF- κ B activation. *Inflamm. Res.* 64, 993–1003
184. Juan, S-H. et al. (2005) Mechanism of concentration-dependent induction of heme oxygenase-1 by resveratrol in human aortic smooth muscle cells. *Biochem. Pharmacol.* 69, 41–48
185. Negi, G. et al. (2011) Nrf2 and NF- κ B modulation by sulforaphane counteracts multiple manifestations of diabetic neuropathy in rats and high glucose-induced changes. *Curr. Neurovasc. Res.* 8, 294–304
186. Yang, Y. et al. (2016) Resveratrol suppresses glial activation and alleviates trigeminal neuralgia via activation of AMPK. *J. Neuroinflammation* 13, 84
187. Benson, C. et al. (2015) Voluntary wheel running delays disease onset and reduces pain hypersensitivity in early experimental autoimmune encephalomyelitis (EAE). *Exp. Neurol.* 271, 279–290
188. Navarro, A. et al. (2004) Beneficial effects of moderate exercise on mice aging: survival, behavior, oxidative stress, and mitochondrial electron transfer. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 286, R505–R511
189. Souza, P.S. et al. (2016) Physical exercise attenuates experimental autoimmune encephalomyelitis by inhibiting peripheral immune response and blood-brain barrier disruption. *Mol. Neurobiol.* Published online July 22, 2016. <http://dx.doi.org/10.1007/s12035-016-0014-0>
190. Grace, P.M. et al. (2016) Prior voluntary wheel running attenuates neuropathic pain. *Pain* 157, 2012–2023
191. Deng, J. et al. (2012) Protective role of reactive oxygen species in endotoxin-induced lung inflammation through modulation of IL-10 expression. *J. Immunol.* 188, 5734–5740
192. Thorpe, G.W. et al. (2013) Superoxide radicals have a protective role during H₂O₂ stress. *Mol. Biol. Cell* 24, 2876–2884
193. Blyth, F.M. et al. (2001) Chronic pain in Australia: a prevalence study. *Pain* 89, 127–134
194. Breivik, H. et al. (2006) Survey of chronic pain in Europe: prevalence, impact on daily life, and treatment. *Eur. J. Pain* 10, 287–333
195. Patton, S. and Kurtz, G.W. (1951) 2-Thiobarbituric acid as a reagent for detecting milk fat oxidation. *J. Dairy Sci.* 34, 669–674
196. Bespalov, I.A. et al. (1999) Fab specific for 8-oxoguanine: control of DNA binding. *J. Mol. Biol.* 293, 1085–1095
197. Souttanakis, R.P. et al. (2000) Fluorescence detection of 8-oxoguanine in nuclear and mitochondrial DNA of cultured cells using a recombinant Fab and confocal scanning laser microscopy. *Free Radic. Biol. Med.* 28, 987–998
198. Griess, P. (1879) Bemerkungen zu der Abhandlung der HH. Weselsky und Benedikt ‘Ueber einige Azoverbindungen’. *Berichte Dtsch. Chem. Ges.* 12, 426–428
199. Feng, L. et al. (2015) P2X7R blockade prevents NLRP3 inflammasome activation and brain injury in a rat model of intracerebral hemorrhage: involvement of peroxynitrite. *J. Neuroinflammation* 12, 190
200. Goswami, S. et al. (2014) Nanomolar detection of hypochlorite by a rhodamine-based chiral hydrazide in absolute aqueous media: application in tap water analysis with live-cell imaging. *Anal. Chem.* 86, 6315–6322
201. Li, G. et al. (2015) A mitochondrial targeted two-photon iridium(III) phosphorescent probe for selective detection of hypochlorite in live cells and in vivo. *Biomaterials* 53, 285–295
202. Li, Z. et al. (2015) A rationally designed upconversion nanoprobe for in vivo detection of hydroxyl radical. *J. Am. Chem. Soc.* 137, 11179–11185
203. Zhuang, M. et al. (2014) Ratiometric fluorescence probe for monitoring hydroxyl radical in live cells based on gold nanoclusters. *Anal. Chem.* 86, 1829–1836
204. Hu, J.J. et al. (2015) Fluorescent probe HKSOX-1 for Imaging and detection of endogenous superoxide in live cells and in vivo. *J. Am. Chem. Soc.* 137, 6837–6843
205. Rong, L. et al. (2016) Hydrogen peroxide detection with high specificity in living cells and inflamed tissues. *Regen. Biomater.* Published online June 11 2016. <http://dx.doi.org/10.1093/rb/rbw022>

206. Kim, S. *et al.* (2014) Far-red fluorescence probe for monitoring singlet oxygen during photodynamic therapy. *J. Am. Chem. Soc.* 136, 11707–11715
207. Mao, Z. *et al.* (2016) NIR in, far-red out: developing a two-photon fluorescent probe for tracking nitric oxide in deep tissue. *Chem. Sci.* 7, 5230–5235
208. Sun, X. *et al.* (2016) A water-soluble copper(II) complex for the selective fluorescence detection of nitric oxide/nitroxyl and imaging in living cells. *ChemPlusChem* 81, 30–34
209. Jing, X. *et al.* (2014) Visualization of nitroxyl (HNO) in vivo via a lysosome-targetable near-infrared fluorescent probe. *Chem. Commun.* 50, 14253–14256
210. Liu, P. *et al.* (2015) A near-infrared fluorescent probe for the selective detection of HNO in living cells and in vivo. *Analyst* 140, 4576–4583
211. Zhang, H. *et al.* (2015) A FRET-based ratiometric fluorescent probe for nitroxyl detection in living cells. *ACS Appl. Mater. Interfaces* 7, 5438–5443
212. Sun, X. *et al.* (2014) A water-soluble boronate-based fluorescent probe for the selective detection of peroxynitrite and imaging in living cells. *Chem. Sci.* 5, 3368–3373
213. Fan, Y. and Ai, H. (2016) Development of redox-sensitive red fluorescent proteins for imaging redox dynamics in cellular compartments. *Anal. Bioanal. Chem.* 408, 2901–2911